



## POST-OP INSTRUCTIONS FOR IMPLANT PATIENTS

PATIENT: \_\_\_\_\_ FOLLOW-UP APPOINTMENT: \_\_\_\_\_

Please read the following instructions carefully. Following these instructions is essential to achieve a successful result. Also, following these instructions will help make your post-operative course more comfortable and will help to avert any unnecessary complications.

1. **REST:** After leaving the office, rest and avoid strenuous activities for the remainder of the day and tomorrow. Keep your head elevated the first two nights after the operation.
2. **DISCOMFORT:** Moderate discomfort is expected for the next few days. If necessary, take the prescribed pain medication as directed. If these do not restore comfort, please call the office.
3. **SWELLING:** Expect some swelling. It may take 3-4 days before it reaches a maximum. Bruising may also occur. Immediately after the surgery, ice packs should be applied to the outside of the face over the area of surgery. Apply for 20 minutes, remove for 20 minutes. Repeat this alternating procedure for the first day only.
4. **BLEEDING:** Slight bleeding may continue for up to 24 hours or so. This is not unusual and is no cause for alarm. Rinsing the mouth interferes with clotting of the blood, so do not rinse vigorously the first evening. If excessive bleeding occurs, it may be stopped by biting continuously for 30-40 minutes on a wet roll of gauze dressing (firm compression). Sitting quietly in an upright position will also aid in the control of bleeding. If these measures fail, call the office.
5. **FEVER:** A slight fever is usual for a day or so following any surgical procedure. Fever may be caused by inadequate fluid intake as well. An adult should consume 8-12 glasses of fluid each day. If any antibiotic was prescribed, be sure to take it as directed. If a fever seems prolonged or excessive, do not hesitate to call the office.
6. **DENTURES:** Dentures may not be worn over the surgical area during the first 10 days. After that period, old dentures may not be worn until refitting adjustment (relining) have been made by your dentist. **INSERTION OF DENTURES TOO EARLY MAY JEOPARDIZE SUCCESSFUL HEALING.**
7. **ORAL HYGIENE:** After today, rinse several times daily with warm salty water, especially after eating (**DO NOT RINSE** the mouth vigorously the day of surgery). Specific mouth rinses or other disinfectants should not be used day of surgery. After the first day, brush your teeth normally, but be careful to stay away from the area of the surgery. **DO NOT** touch the surgical area.
8. **DIET:** Only soft food should be consumed for the first 5 days to avoid trauma to the surgical sites. After that, you may advance your diet as tolerated. If you have no teeth, only liquids may be consumed during the first 5 days. Semi-liquid (mashed) food may be added to the diet after the 5th day.
9. **PROBLEMS:** If you have questions or if there is any sign of a surgery related problem, call the office.
10. **ALCOHOL/SMOKING:** **ALCOHOL AND SMOKING ARE ABSOLUTELY PROHIBITED FOR THE FIRST TWO WEEKS.** Alcoholic beverages can close blood vessels and destroy blood clots. Smoking is an irritant to healing tissues and will also close down blood vessels. Both will retard healing.

3929 S. Hwy 14  
Greenville, SC  
864-281-9119

700 Squires Point Suite A  
Duncan, SC  
864-486-8330

107 Clair Dr.  
Powdersville, SC  
864-565-8760

2 Cleveland Court  
Greenville, SC  
864-271-9990